

Sequences with a Unique Realization by Simple Graphs*

MICHAEL KOREN[†]

*Department of Operations Research, College of Engineering,
Cornell University, Ithaca, New York*

Communicated by W. T. Tutte

Received May 3, 1974

Sequences with unique realizations (up to isomorphism) by simple graphs are characterized, partly by reduction to the case of pairs of sequences, which have a unique realization by bipartite graphs (see Theorem 6.1).

1. INTRODUCTION

All graphs in this paper are finite, undirected, and without loops or multiple edges.

$G(p_1, \dots, p_n)$ denotes a graph with n vertices p_1, \dots, p_n . If two vertices p, q are adjacent in G , we write $(p, q) \in G$.

$d(p_i)$ denotes the degree of p_i in G .

A sequence $\psi = (b_1, \dots, b_n)$ is *graphic* if there exists a graph $G = G(p_1, \dots, p_n)$ such that $d(p_i) = b_i$ for $i = 1, \dots, n$. G is called a *realization* of ψ . We say also that G *realizes* ψ . If every two realizations of ψ are isomorphic we call ψ *unigraphic*.

Similarly, a pair of sequences $[(b_1, \dots, b_n), (c_1, \dots, c_m)]$ is *graphic* if $d(p_i) = b_i$ ($i = 1, \dots, n$) and $d(q_i) = c_i$ ($i = 1, \dots, m$) for some bipartite graph $G(p_1, \dots, p_n, q_1, \dots, q_m)$, all its edges are of the form (p_i, q_j) .

A graphic pair is *unigraphic* if any two realizations of it are isomorphic.

In this paper we characterize unigraphic sequences. One of the criteria for unigraphicness of a sequence is the unigraphicness of a pair of sequences. A characterization of unigraphic pairs was given in [5].

* Supported by the Office of Naval Research under Contract N00014-67-A-0077-0014.

[†] Present address, Department of Mathematics, Ben Gurion University, Beer-Sheva, Israel.

By acceptance of this article, the publisher or recipient acknowledges the U. S. Government's right to retain a nonexclusive, royalty-free license in and to any copy-right covering the article.

In [2], Erdős and Gallai show that a nonincreasing sequence (b_1, \dots, b_n) of natural numbers is graphic iff:

$$\sum_{i=1}^n b_i \text{ is even} \quad (1.1)$$

$$\sum_{i=1}^j b_i - j(j-1) \leq \sum_{i=j+1}^n \min(j, b_i), \quad (j = 1, \dots, n). \quad (1.2)$$

Notation. Throughout this paper ϕ will denote the graphic non-increasing sequence (a_1, \dots, a_n) of positive integers, and a , with some lower index, will denote a term of ϕ .

DEFINITION 1.1. For a nonincreasing sequence $\psi = (b_1, \dots, b_m)$ we define

$$\delta(j, \psi) = \sum_{i=j+1}^m \min(j, b_i) - \sum_{i=1}^j b_i + j(j-1) \quad (\text{for } j = 0, 1, \dots, m).$$

Remark. Of course, $\delta(0, \psi) = 0$ for any ψ . $\delta(j, \phi) \geq 0$ for $j = 1, \dots, n$; $\delta(n, \phi) = 0$ iff $a_1 = a_n = n - 1$, otherwise $\delta(n, \phi) > 1$.

As always, $K = K_n$ denotes the complete graph on n vertices. $K^c = K_n^c$ denotes the empty graph on n vertices; i.e., the graph with no edges.

Similarly, $K = K_{n,m}$ denotes the complete bipartite graph with two independent sets having n and m vertices, respectively. $K^c = K_{n,m}^c$ denotes the empty graph on $n + m$ vertices.

If X and Y are sets of vertices of G , $G[X, Y]$ denotes the subgraph of G , generated by X, Y , i.e., the graph with $X \cup Y$ as its set of vertices, which includes exactly those edges of G having one endpoint in X and one endpoint in Y . We write $G[X]$ for $G[X, X]$.

2. THE STRUCTURE OF SOME GRAPHS

LEMMA 2.1. Suppose $\delta(j, \phi) = 0$ for some j , $1 \leq j < n$. If $a_{j+1} > j$, let k be an index such that $a_k \geq j \geq a_{k+1}$. If $a_{j+1} \leq j$, let $k = j$. For any realization $G = G\{p_1, \dots, p_n\}$ of ϕ , define $S = \{p_1, \dots, p_j\}$, $T = \{p_{k+1}, \dots, p_n\}$, $U = \{p_{j+1}, \dots, p_k\}$. Then:

$$G[S] = K \quad (2.1)$$

$$G[T] = K^c \quad (2.2)$$

and if $U \neq \emptyset$

$$G[S, U] = K \quad (2.3)$$

$$G[T, U] = K^c. \quad (2.4)$$

Proof. $\sum_{i=1}^j a_i - j(j-1)$ and $\sum_{i=j+1}^n \min(j, a_i)$ are lower and upper bounds, respectively, on the number of edges of $G[S, T \cup U]$; since they are equal there must be $\frac{1}{2}j(j-1)$ edges in $G[S]$, and $\min(j, a_i)$ edges in $G[S, \{p_i\}]$ for $i > j$; and the results follow (see Fig. 1).




FIG. 1. The structure of G , when $\delta(j, \phi) = 0$.

The converse Lemma is also true:

LEMMA 2.2. Suppose $G(p_1, \dots, p_n)$ realizes ϕ , $S = \{p_1, \dots, p_j\}$, $n > j \geq 1$, $T = \{p_{k+1}, \dots, p_n\}$, $k \geq j$, $U = \{p_{j+1}, \dots, p_k\}$ and conditions (2.1), (2.2) hold for S and T and, if $U \neq \emptyset$, conditions (2.3), (2.4) hold as well. Then $\delta(j, \phi) = 0$.

Proof. The proof is straightforward and is omitted (see Fig. 1).

DEFINITION 2.1. $h = h(j) = \max\{j, \max\{i: a_i > j\}\}$; i.e., $h = j$ if $a_{j+1} \leq j$, $h = n$ if $a_n > j$, and if $a_{j+1} > j \geq a_n$, then h is defined by $a_h > j \geq a_{h+1}$.

LEMMA 2.3. Suppose $a_1 = a_2 = \dots = a_l > a_{l+1} \geq \dots \geq a_n$ ($1 \leq l \leq n$), $\delta(j, \phi) = 1$ for some j , $1 \leq j < l$ and $j \geq a_n$. Then $\delta(j+1, \phi) \leq 1$.

Proof. As $1 + \sum_{i=1}^j a_i - j(j-1) = \sum_{i=j+1}^n \min(j, a_i)$, we must have $a_{j+1} > j$, since otherwise

$$1 + \sum_{i=1}^j a_i - j(j-1) = \sum_{i=j+1}^n a_i,$$

a contradiction of condition (1.1). Hence $n > h > j$. Thus

$$1 + \sum_{i=1}^j a_i = j(j-1) + j(h-j) + \sum_{i=h+1}^n a_i. \quad (2.5)$$

Hence

$$ja_1 \geq j(h-1), \quad \text{or} \quad a_1 \geq h-1.$$

Suppose

$$1 + \sum_{i=1}^{j+1} a_i < (j+1)j + (j+1)(h-j-1) + \sum_{i=h+1}^n a_i. \quad (2.6)$$

Subtracting (2.5) from (2.6) we obtain $a_1 = a_{j+1} < h-1$, a contradiction.

LEMMA 2.4. *Suppose $\delta(j, \phi) = 0$ for some j , $1 \leq j < n$, then ϕ is unigraphic iff the pair $[(a_1 - (h-1), \dots, a_j - (h-1)), (a_{h+1}, \dots, a_n)]$ is unigraphic and so is the sequence $(a_{j+1} - j, \dots, a_h - j)$, (if $h > j$).*

Proof. Lemma 2.4 follows easily from Lemma 2.1.

3. PROPERTIES OF UNIGRAPHIC SEQUENCES

We will need the following known Theorem:

THEOREM 3.1. ([2] and compare [3].) *If $\psi = (b_1, \dots, b_n)$ is graphic, then for each j , $j = 1, \dots, n$, there exists a realization $G_j(p_1, \dots, p_n)$ of ψ in which p_j is adjacent to the first (other) b_j vertices with the highest degrees.*

COROLLARY 3.1. *If $a_1 > a_2 > a_{n-1} > a_n$, then ϕ has a realization $G = G(p_1, \dots, p_n)$, such that $(p_{n-1}, p_1), (p_2, p_1) \in G$ and $(p_{n-1}, p_n), (p_2, p_n) \notin G$.*

Proof. By Theorem 3.1, $\psi^* = (a_1 - 1, a_3 - 1, \dots, a_{a_2+1} - 1, a_{a_2+2}, \dots, a_n)$ is graphic. By the same theorem ψ^* has a realization $G_{n-1}(p_1, p_3, \dots, p_n)$, in which $(p_{n-1}, p_1) \in G_{n-1}$ and $(p_{n-1}, p_n) \notin G_{n-1}$. Adding the vertex p_2 and connecting it to $p_1, p_3, p_4, \dots, p_{a_2+1}$ we obtain the desired realization G .

DEFINITION 3.1. *If $(p_{i_1}, p_{i_2}) \in G$ ($(p_{i_1}, p_{i_2}) \notin G$) then $G - (p_{i_1}, p_{i_2})$ ($G + (p_{i_1}, p_{i_2})$) denotes the graph which is obtained from G by removing (adding) the edge (p_{i_1}, p_{i_2}) .*

The next Theorem is one of our main results.

THEOREM 3.2. *If $a_2 > a_{n-1}$ ($n \geq 4$) and ϕ is unigraphic, then $\delta(j, \phi) = 0$ for some j , $1 \leq j < n$.*

Proof. Assume $\delta(j, \phi) > 0$ for $j = 1, \dots, n-1$. We will consider two cases.

Case a. $\delta(j, \phi) > 1$ for $j = 1, \dots, n-1$.

In this case $\phi^* = (a_1 + 1, a_2, \dots, a_{n-1}, a_n - 1)$ is graphic and, by Corollary 3.1, ϕ^* has a realization $G^* = G^*(p_1, \dots, p_n)$ such that $(p_2, p_1), (p_{n-1}, p_1) \in G^*$ and $(p_2, p_n), (p_{n-1}, p_n) \notin G^*$. Thus $G_1 = G^* - (p_2, p_1) + (p_1, p_n)$ and $G_2 = G^* - (p_{n-1}, p_1) + (p_{n-1}, p_n)$ realize ϕ . Furthermore, G_1 and G_2 are not isomorphic since the number of edges between a vertex of degree a_1 and a vertex of degree a_2 is smaller in G_1 than in G_2 .

Case b. $\delta(j, \phi) = 1$ for some values of j .

Let j_0 be the largest such j (i.e., $\delta(j_0, \phi) = 1$ and $\delta(j, \phi) > 1$ for $j > j_0$).

Since $\delta(j_0, \phi) = 1$ we have, by condition (1.1), $a_{j_0+1} > j_0$. If $h = h(j_0) = n$ (see Definition 2.1), then we proceed as in case a. Assume, therefore, that $n > h > j_0$. Then the sequence $\psi = (a_1 + 1, a_2, \dots, a_{h-1}, a_h - 1, a_{h+1}, \dots, a_n)$ is graphic and $\delta(j_0, \psi) = 0$. Since $a_n \leq j_0 < h$, ψ has, by Theorem 3.1, a realization $G' = G'(p_1, \dots, p_n)$ such that $(p_n, p_1) \in G'$ and $(p_n, p_h) \notin G'$. Therefore $G = G' - (p_1, p_n) + (p_h, p_n)$ realizes ϕ and for $S = \{p_1, \dots, p_{j_0}\}$, $T = \{p_{h+1}, \dots, p_n\}$, $U = \{p_{j_0+1}, \dots, p_h\}$; conditions (2.1) (2.2) and (2.3) hold, and $G[T, U] = K^c + (p_h, p_n)$.

If $h = j_0 + 1$, then, by Lemma 2.2, $\delta(j_0 + 1, \phi) = 0$, a contradiction. Hence $h > j_0 + 1$.

Since $G[T, U - \{p_h\}] = K^c$, we have

$$h - 1 \geq d(p_{j_0+1}) = a_{j_0+1} \geq a_h = d(p_h).$$

But $G[S, U] = K$, hence there exists an index r , $h > r > j_0$, such that $(p_r, p_h) \notin G$. Of course, $(p_r, p_1) \in G$, and thus another realization of ϕ is $G_1 = G' - (p_1, p_r) - (p_h, p_n) + (p_1, p_n) + (p_r, p_h)$. But $d(p_r) > d(p_n)$ and hence by the unigraphicity of ϕ we have $d(p_1) = d(p_h)$. Thus, by Lemma 2.3, we arrived at a contradiction to the maximality of j_0 .

The next theorem sharpens the results of Theorem 3.2:

THEOREM 3.3. *Suppose ϕ is unigraphic, j_0 is the largest index j for which $\delta(j, \phi) = 0$ and $h' = h(j_0) > j_0$. Then $h' \geq j_0 + 3$, $a_{j_0+1} < h' - 1$ and $a_{j_0+2} = a_{h'-1}$.*

Proof. By Lemma 2.4, the sequence $\psi = (a_{j_0+1} - j_0, \dots, a_{h'} - j_0) = (b_1, \dots, b_m)$ is unigraphic. $b_m > 0$, hence $m \geq 2$. If $m = 2$, then $\psi = (1, 1)$ and if $m = 3$, then $\psi = (2, 1, 1)$ or $\psi = (2, 2, 2)$. In each case $\delta(1, \psi) = 0$.

Suppose $m \geq 4$ and $b_2 > b_{m-1}$. Then, by Theorem 3.2, there exists j , $1 \leq j < m$, such that $\delta(j, \psi) = 0$.

To complete the proof of the Theorem it is enough to show that the assumption $\delta(j, \psi) = 0$ for some j , $1 \leq j < m$ leads to a contradiction.

Suppose that $\delta(j, \psi) = 0$ for some j , $1 \leq j < m$. Fix such j . Let $G(p_1, \dots, p_n)$ be a realization of ϕ . Define $S_1 = \{p_1, \dots, p_{j_0}\}$, $T_1 = \{p_{h'+1}, \dots, p_n\}$, $U_1 = \{p_{j_0+1}, \dots, p_h\}$. S_1 , T_1 , U_1 fulfill conditions (2.1) to (2.4), by Lemma 2.1. Define $h'' = h(j)$ (in ψ), $S_2 = \{p_{j_0+1}, \dots, p_{j_0+j}\}$, $T_2 = \{p_{j_0+h''+1}, \dots, p_h\}$, $U_2 = \{p_{j_0+j+1}, \dots, p_{j_0+h''}\}$. By Lemma 2.1, S_2 , T_2 , U_2 (in $G[U_1]$) fulfill conditions (2.1) to (2.4). Define $S = S_1 \cup S_2$, $T = T_1 \cup T_2$, $U = U_2$. It is easy to check that S , T and U also fulfill conditions (2.1) to (2.4). Hence, by Lemma 2.2, $\delta(j_0 + j, \phi) = 0$, a contradiction to the maximality of j_0 .

Up to now we assumed always $a_2 > a_{n-1}$. In the next two sections we will abandon this assumption.

4. THE REGULAR CASE

DEFINITION 4.1. ϕ^c , the complementary sequence of ϕ , is defined by $\phi^c = (a_1^c, \dots, a_n^c) = (n-1-a_1, \dots, n-1-a_n)$.

Remark. The complementary graph of any realization of ϕ , is a realization of ϕ^c . Hence ϕ is unigraphic iff ϕ^c is unigraphic.

THEOREM 4.1. Suppose $a_1 = a_n$. Then ϕ is unigraphic iff $a_1 \in \{1, n-2, n-1\}$ or $\phi = (2, 2, 2, 2, 2)$.

Proof. We may assume $a_1 \leq (n-1)/2$, since otherwise $a_1^c < (n-1)/2$.

Case a. $1 < a_1 < (n-1)/2$. Let $\phi_1 = (a_1, \dots, a_{a_1+1})$, $\phi_2 = (a_{a_1+2}, \dots, a_n)$. Both ϕ_1 and ϕ_2 are graphic and hence ϕ has a disconnected realization. On the other hand, $\sum_{i=1}^n a_i = na_1 > 2(n-1)$ and therefore ϕ has a connected realization (see [1; 2, (1.6)]).

Case b. $a = (n-1)/2$. In this case n is odd, hence a is even. Let $a = 2m$. Choose a polytope on n vertices, p_1, \dots, p_n (in this order). By adding to the polytope all the diagonals with "length" $< m$ we obtain a graph $G = G(p_1, \dots, p_n)$ which realizes ϕ . (Where the length of a diagonal is the number of edges of the polytope, on the shortest path between the endpoints of the diagonal). No set of $m+2$ vertices form in G a clique (i.e., for no set $Y = \{p_{i_1}, \dots, p_{i_{m+2}}\}$, does $G[Y]$ equal K_{m+2}).

Let $X = \{p_2, p_3, \dots, p_{m+3}\}$. $G[X] = K_{m+2} - (p_2, p_{m+3})$. If $a_1 > 2$ ($m > 1, n > 9$), then $(p_1, p_{m+4}) \notin G$. Hence $G^* = G - (p_1, p_2) - (p_{m+3}, p_{m+4}) + (p_1, p_{m+4}) + (p_2, p_{m+3})$ realizes ϕ . G^* is not isomorphic to G since $G^*[X] = K_{m+2}$.

5. THE ALMOST REGULAR CASES

LEMMA 5.1. *If $n - 1 > a_1 > a_2 \geq \dots \geq a_{n-1} > a_n$, then ϕ is not unigraphic.*

Proof. Using Theorem 3.1 twice we obtain a realization $G_1(p_1, \dots, p_n)$ of ϕ and a realization $G_n(p_1, \dots, p_n)$ of ϕ . In G_1 , p_1 is adjacent to $p_2, p_3, \dots, p_{a_1+1}$, and hence p_1 is not adjacent to p_n . In G_n , p_n is adjacent to p_1, p_2, \dots, p_{a_n} . Therefore G_1 and G_n are not isomorphic.

LEMMA 5.2. (a) *If $a_1 > a_2 = a_n$, then ϕ is unigraphic iff $(a_2 - 1, a_3 - 1, \dots, a_{a_1+1} - 1, a_{a_1+2}, \dots, a_n)$ is unigraphic.*

(b) *If $a_1 = a_{n-1} > a_n$, then ϕ is unigraphic iff $(a_1 - 1, \dots, a_{a_n} - 1, a_{a_n+1}, \dots, a_{n-1})$ is unigraphic.*

Proof. The Lemma is self-evident.

In light of Lemma 5.2 we will investigate, in the next two lemmas, sequences of the form $(a, \dots, a, a - 1, \dots, a - 1)$.

LEMMA 5.3. *If $n - 1 > a_1 = a_k = a_{k+1} + 1 = a_n + 1$ and $2 \leq k \leq n - 2$, then*

(a) $\delta(j, \phi) > 0$ for $j \neq k$.

(b) $\delta(k, \phi) = 0$ iff $\phi = (2, 2, 1, 1)$.

Proof. If $\delta(j, \phi) = 0$ for some j ($1 \leq j < n$), then $a_n < j$, since otherwise:

$$\begin{aligned} j(n - j) &= j(n - 1) - j(j - 1) > \sum_{i=1}^j a_i - j(j - 1) \\ &= \sum_{i=j+1}^n \min(j, a_i) = j(n - j), \quad \text{a contradiction.} \end{aligned}$$

But $a_1 > j - 1$, since otherwise

$$0 \geq \sum_{i=1}^j a_i - j(j - 1) = \sum_{i=j+1}^n \min(j, a_i) \geq a_n > 0, \quad \text{a contradiction.}$$

Hence $j \geq a_1 > j - 1$, or $a_1 = j$.

Suppose $\delta(j, \phi) = 0$ for $j < k$. Then

$$\begin{aligned} j &= j^2 - j(j - 1) = \sum_{i=1}^j a_i - j(j - 1) = \sum_{i=j+1}^n \min(j, a_i) \\ &= j(k - j) + (j - 1)(n - k) > j, \quad \text{a contradiction.} \end{aligned}$$

($j - 1 = a_1 - 1 \geq 1, n - k > 1$).

Suppose $\delta(j, \phi) = 0$ for $n > j > k$. Then $k = \sum_{i=1}^j a_i - j(j-1) = (j-1)(n-j) \geq j-1 \geq k$. Hence $n-j=1$, or $n-a_1=1$, a contradiction.

Suppose $\delta(k, \phi) = 0$. Then

$$k = \sum_{i=1}^k a_i - k(k-1) = (k-1)(n-k). \quad (5.1)$$

$n-k \geq 2$, hence $k \geq 2(k-1)$, or $2 \geq k$. Since $k \leq 2$ we have $a_1 = k = 2$ and from condition (5.1), $n = 4$.

LEMMA 5.4. *If $\psi = (b, \dots, b, b-1, \dots, b-1)$, where b occurs $k \geq 2$ times and $b-1$ occurs $n-k \geq 2$ times, then ψ is unigraphic iff one of the following holds:*

- (a) $\psi = (2, 2, 1, 1)$.
- (b) $\psi = (1, \dots, 1, 0, \dots, 0)$ (and k is even).
- (c) $\psi = (n-1, \dots, n-1, n-2, \dots, n-2)$ (and $n-k$ is even).

Proof. If $n-1 > b > 1$, then, by Theorem 3.2, ψ is unigraphic only if $\delta(j, \psi) = 0$ for some $1 \leq j < n$. By Lemma 5.3, $\delta(j, \psi) = 0$ only if $\psi = (2, 2, 1, 1)$ (and $j = 2$). The sequence $(2, 2, 1, 1)$ is of course unigraphic. If $b = 1$ then ψ is unigraphic for any even k . If $b = n-1$, then $\psi^c = (0, \dots, 0, 1, \dots, 1)$ and hence ψ^c and ψ are unigraphic for any even $n-k$.

From Lemmas 5.1 to 5.4 we obtain, in the next Lemma, a characterization for unigraphic sequences in which $a_2 = a_{n-1}$.

LEMMA 5.5. *Let $\psi = (b_1, \dots, b_n)$ be a graphic sequence, $n-1 > b_i > 0$ ($i = 1, \dots, n$) and $b_1 \neq b_2 = b_3 = \dots = b_n$. Then ψ is unigraphic iff one of the following conditions holds:*

- (1) $b_2 = 1$
- (2) $b_1 = n-2$ and $b_2 = 2$
- (3) $b_2 = n-2$
- (4) $b_1 = 1$ and $b_2 = n-3$

Proof. If condition (3) or condition (4) holds for ψ , then condition (1) or condition (2), respectively, holds for ψ^c . Hence we may assume that $b_1 > b_2$.

By Lemma 5.2, ψ is unigraphic iff $\psi_1 = (b_2, \dots, b_k, b_{k+1}-1, \dots, b_n-1)$ is unigraphic, where $k = n - b_1$. $n-1 > b_1 > 1$, hence $n-2 \geq k \geq 2$.

Case a. $b_1 < n - 2$. In this case $k > 2$ and we may apply Lemma 5.4 to ψ_1 . $\psi_1 \neq (2, 2, 1, 1)$. Since otherwise $\psi = (2, \dots, 2)$. $n - 1 > b_1 > b_2$, hence $b_2 < (n - 1) - 1$; thus $\psi_1 \neq ((n - 1) - 1, \dots, (n - 1) - 1, (n - 1) - 2, \dots, (n - 1) - 2)$. Therefore ψ_1 is unigraphic iff $\psi_1 = (1, \dots, 1, 0, \dots, 0)$ and $\psi = (b_1, 1, \dots, 1)$.

Case b. $b_1 = n - 2$, or $k = 2$. In this case $\psi_1 = (b_2, b_3 - 1, \dots, b_n - 1)$. $b_2 < (n - 1) - 2$, since the sequence $(n - 2, n - 3, \dots, n - 3)$ is not graphic. Thus we may apply case a to ψ_1 . Therefore $\psi_1 = (b_2, 1, \dots, 1)$. Hence $b_2 = b_3 = 2$ and $\psi = (n - 2, 2, \dots, 2)$.

6. CONCLUSIONS

In this short section we summarize our results into one main theorem:

THEOREM 6.1 (MAIN THEOREM). *Given a sequence ϕ , $n > 3$, let j_0 be the largest index j , such that $\delta(j, \phi) = 0$, ($0 \leq j \leq n$); and let $h = h(j_0)$ (see Definition 2.1).*

If $a_2 = a_{n-1}$, then ϕ is unigraphic if and only if one of the following conditions holds:

$$a_1 = a_n, \quad a_n \in \{1, n - 2, n - 1\} \quad (6.1)$$

$$a_1 = a_n = 2, \quad n = 5 \quad (6.2)$$

$$a_1 > a_2 = a_n = 1 \quad (6.3)$$

$$a_1 > a_2 = a_n = 2, \quad a_1 \in \{n - 1, n - 2\} \quad (6.4)$$

$$n - 2 = a_1 = a_{n-1} > a_n \quad (6.5)$$

$$n - 3 = a_1 = a_{n-1} > a_n = 1 \quad (6.6)$$

$$n - 1 = a_1 > a_2 = a_n = 3, \quad n = 6 \quad (6.7)$$

$$n - 1 = a_1 > a_2 = a_n = n - 2. \quad (6.8)$$

If $a_2 > a_{n-1}$, then ϕ is unigraphic if and only if

(1) *either $h = n$, or the pair $[(a_1 - (h - 1), \dots, a_{j_0} - (h - 1)), (a_{h+1}, \dots, a_n)]$ is unigraphic, and*

(2) *either $h = j_0$, or $h > j_0 + 3$ and the sequence $(a'_1, \dots, a'_n) = (a_{j_0+1} - j_0, \dots, a_n - j_0)$ fulfills one of the conditions (6.1)-(6.6).*

Proof. The Theorem follows easily from Lemma 2.4, Theorem 3.2, Theorem 3.3, Theorem 4.1, Theorem 5.1, and Lemma 5.5.

Remark. While writing this paper, the author learned that Shuo-Yen R. Li wrote a paper on the same subject [6]. In particular Theorem 4.1 and Lemma 5.2 appear in [6] too. However, the attitude and the nature of his results are different from ours.

REFERENCES

1. J. EDMONDS, Existence of k -edge connected ordinary graphs with prescribed degrees, *J. Res. Nat. Bur. Stand. Sect. B* **68B** (1964), 73–74.
2. P. ERDÖS AND T. GALLAI, Gráfok elöirt fokú pontokkal, *Mat. Lapok* **11** (1960), 264–274.
3. S. L. HAKIMI, On the realizability of a set of integers as degrees of the vertices of a linear graph I, *J. SIAM* **10** (1962), 496–506.
4. F. HARARY, “Graph theory.” Addison-Wesley, Reading, 1969.
5. M. KOREN, Pairs of sequences with a unique realization by bipartite graphs, *J. Combinatorial Theory (B)*, this issue.
6. SHUO-YEN R. LI, Graphic sequences with unique realization, *J. Combinatorial Theory (B)* **19** (1975), 42–68.